

prosci-inc.com






HIGH PERFORMANCE ANTIBODIES ... AND MORE

ProSci Incorporated 12170 Flint Place Poway, CA 92064 Toll Free: +1 (888) 513 9525 Local: +1 (858) 513 2638 Fax: +1 (858) 513 2692


techsupport@prosci-inc.com

ZFP1 Antibody

CATALOG NUMBER: 27-369



Antibody used in IHC on Human Spermatophore at 4.0-8.0 ug/ml.



Antibody used in IHC on Human Intestine at 4.0-8.0 ug/ml.



Antibody used in WB on Human Jurkat 0.2-1 ug/ml.

Specifications	
SPECIES REACTIVITY:	Human
TESTED APPLICATIONS:	ELISA, IHC, WB
APPLICATIONS:	ZFP1 antibody can be used for detection of ZFP1 by ELISA at 1:62500. ZFP1 antibody can be used for detection of ZFP1 by western blot at 0.1-2.0 ug/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000.
USER NOTE:	Optimal dilutions for each application to be determined by the researcher.
POSITIVE CONTROL:	1) Cat. No. 1205 - Jurkat Cell Lysate
PREDICTED MOLECULAR WEIGHT:	41 kDa
IMMUNOGEN:	Antibody produced in rabbits immunized with a synthetic peptide corresponding a region of human ZFP1.
HOST SPECIES:	Rabbit
Properties	
PURIFICATION:	Antibody is purified by peptide affinity chromatography method.
PHYSICAL STATE:	Lyophilized
BUFFER:	Antibody is lyophilized in PBS buffer with 2% sucrose. Add 50 uL of distilled water. Final antibody concentration is 1 mg/mL.
CONCENTRATION.	1

PHYSICAL STATE:	Lyophilized
BUFFER:	Antibody is lyophilized in PBS buffer with 2% sucrose. Add 50 uL of distilled water. Final antibody concentration is 1 mg/mL.
CONCENTRATION:	1 mg/ml
STORAGE CONDITIONS:	For short periods of storage (days) store at 4°C. For longer periods of storage, store ZFP1 antibody at -20°C. As with any antibody avoid repeat freeze-thaw cycles.
CLONALITY:	Polyclonal
CONJUGATE:	Unconjugated

Additional Info	
ALTERNATE NAMES:	ZFP1, ZNF475
ACCESSION NO.:	NP_710155
PROTEIN GI NO.:	119226229

OFFICIAL SYMBOL:	ZFP1
GENE ID:	162239
Background	
BACKGROUND:	Zinc finger proteins (Zfp) are encoded by a large family of genes present in many organisms including yeast and human. Some of them are transcriptional activators and bind specifically to DNA by zinc mediated folded structures commonly known as zinc fingers. The putative Zfp-1 (Zinc finger protein 1 homolog) protein contains in addition to 7 zinc fingers, two helix-turn-helix motifs.
REFERENCES:	1) Chowdhury, K., et al., (1989) NucleicAcidsRes.17(24), 10427-10438.

FOR RESEARCH USE ONLY

December 12, 2016